NEW DRUG REVIEWS FOR 06/21/2012 PA MEETING		
OPEN ACCESS		
COMMON TRADE NAME	GENERIC NAME	INDICATIONS
Diazepam 5mg/5ml cup Solution	Diazepam	Indicated for the management of anxiety disorders or for the short-term relief of the symptoms of anxiety. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic.
Cosopt PF 2%-0.5% Droperette	DORZOLAMIDE/TIMOLOL/PF	Indicated for the reduction of elevated intraocular pressure (IOP) in patients with open-angle glaucoma or ocular hypertension who are insufficiently responsive to beta-blockers (failed to achieve target IOP determined after multiple measurements over time).
Picato 0.05% Gel (EA)	INGENOL MEBUTATE	An inducer of cell death indicated for the topical treatment of actinic keratosis.
Picato 0.015% Gel (EA)	INGENOL MEBUTATE	An inducer of cell death indicated for the topical treatment of actinic keratosis.
Hydrocodone Acetaminophen 7.5-325/CP Solution	HYDROCODONE BIT/ACETAMINOPHEN	Indicated for the relief of moderate to moderately severe pain.
Naftin 2% Cream (G)	NAFTIFINE HCL	Used for skin infections such as athlete's foot, jock itch, and ringworm.
Kalydeco 150mg Tablet	IVACAFTOR	Indicated for the treatment of a rare form of cystic fibrosis (CF) in patients ages 6 years and older who have the specific G551D mutation in the Cystic Fibrosis Transmembrane Regulator (CFTR) gene.
Hylase 2.5% Gel	HYALURONATE SODIUM/HE-CELL/PEG	Indicated for the management and treatment of leg ulcers, pressure ulcers, diabetic ulcers, and exuding wounds prone to bleeding.
Xyntha Solofuse 250(+/-) Syringe Kit	ANTIHEMOPH.FVIII PLAS/ALB FREE	Indicated for the control and prevention of bleeding episodes in patients with hemophilia A (congenital factor VIII deficiency or classic hemophilia) and for surgical prophylaxis in patients with hemophilia A.
Xyntha Solofuse 500(+/-) Syringe Kit	ANTIHEMOPH.FVIII PLAS/ALB FREE	Indicated for the control and prevention of bleeding episodes in patients with hemophilia A (congenital factor VIII deficiency or classic hemophilia) and for surgical prophylaxis in patients with hemophilia A.
Prenata 29mg-1mg Tab Chewable	PRENATAL VIT37/IRON/FOLIC ACID	Prenatal Vitamin
Rectiv 0.4% Ointment (G)	NITROGLYCERIN	Indicated for the treatment of moderate to severe pain associated with chronic anal fissures.
Promacta 12.5mg Tablet	Eltrombopag Olamine	Indicated for the treatment of thrombocytopenia in patients with chronic immune (idiopathic) thrombocytopenic purpura (ITP) who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy.
Citranatal Harmony 30-1-50mg Capsule	PNV34/IRON,CARBONYL/FA/DSS/DHA	Prenatal Vitamin
B-Nexa 40-1.2-100 Tablet	VIT B6/FA/CALCIUM PHOS/GINGER	Prenatal Vitamin

PDL EDITS FOR IMPLEMENTATION		
COMMON TRADE NAME	GENERIC NAME	INDICATIONS
Intermezzo 3.5mg Tablet Sublingual	ZOLPIDEM TARTRATE	Sedative Hypnotics PDL Edit Non-Preferred
Intermezzo 1.75mg Tablet Sublingual	ZOLPIDEM TARTRATE	Sedative Hypnotics PDL Edit Non-Preferred
Zioptan 0.0015% Droperette	TAFLUPROST/PF	Ophthalmic Prostaglandin Agonists PDL Edit Non Preferred
Janumet XR 100-1000mg TBMP Tablet	SITAGLIPTIN PHOS/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Janumet XR 50-1000mg TBMP Tablet	SITAGLIPTIN PHOS/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Janumet XR 50-500mg TBMP Tablet	SITAGLIPTIN PHOS/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Bydureon 2mg Vial	EXENATIDE MICROSPHERES	GLP-1 Receptor Agonists PDL Edit Non-Preferred
Jentadueto 2.5mg-500mg Tablet	LINAGLIPTIN/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Jentadueto 2.5mg-850mg Tablet	LINAGLIPTIN/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Jentadueto 2.5mg-1000mg Tablet	LINAGLIPTIN/METFORMIN HCL	Oral Anti-Diabetics Combinations PDL Edit Preferred
Edarbyclor 40-25mg Tablet	AZILSARTAN MED/CHLORTHALIDONE	Angiotensin II Receptor Blocker Diuretic Combinations
		Non-Preferred
Edarbyclor 40-12.5mg Tablet	AZILSARTAN MED/CHLORTHALIDONE	Angiotensin II Receptor Blocker Diuretic Combinations
		Non-Preferred
Pedipirox-4 8% Kit	CICLOPIROX/NAIL LACQ/FT DEOD#4	Onychomycosis Antifungal Agents PDL Edit Non-Preferred
Dutoprol 100-12.5mg Tablet ER	METOPROLOL SUCCINATE/HCTZ	Beta Adrenergic Blockers PDL Edit <b>Preferred</b>
Dutoprol 50-12.5mg Tablet ER	METOPROLOL SUCCINATE/HCTZ	Beta Adrenergic Blockers PDL Edit Preferred
Dutoprol 25-12.5mg Tablet ER	METOPROLOL SUCCINATE/HCTZ	Beta Adrenergic Blockers PDL Edit Preferred
CLINICAL EDITS FOR IMPLEMENTATION		
COMMON TRADE NAME	GENERIC NAME	INDICATIONS
Relistor 8mg/0.4mL Disp Syringe	Methylnaltrexone Bromide	Indicated for the treatment of opioid-induced constipation in patients
	'	with advanced illness who are receiving palliative care, when response
		to laxative therapy has not been sufficient
Relistor 12mg/0.6mL Disp Syringe	Methylnaltrexone Bromide	Indicated for the treatment of opioid-induced constipation in patients
	,	with advanced illness who are receiving palliative care, when response
		to laxative therapy has not been sufficient
Latuda 20mg Tablet	LURASIDONE HCL	Atypical Antipsychotics Clinical Edit
Oxecta 5mg Tablet	OXYCODONE HCL	Short Acting Narcotics Clinical Edit
Oxecta 7.5mg Tablet	OXYCODONE HCL	Short Acting Narcotics Clinical Edit
REQUEST FOR CONTINUED PRIOR		
COMMON TRADE NAME		
Lidocaine 5% Ointment (G)	Lidocaine	Indicated for production of anesthesia of accessible mucous
	Lidodinio	membranes of the oropharynx. It is also useful as an anesthetic
		lubricant for intubation and for the temporary relief of pain associated
		with minor burns, including sunburn, abrasions of the skin, and insect
		bites.
Aqua Glycolic HC 2% Combo Package	HYDROCORTISONE/SKIN CLNSR #25	Used to treat or prevent dry skin.
Orbivan CF 50mg-300mg Tablet	BUTALBITAL/ACETAMINOPHEN	Indicated for the relief of the symptom complex of tension (or muscle
		contraction) headache.
Zyclara 3.75% Cream	Imiquimod	Used to treat certain types of actinic keratoses (flat, scaly growths on
	•	the skin caused by too much sun exposure) on the face or scalp.